



Calhoun: The NPS Institutional Archive

Reports and Technical Reports

All Technical Reports Collection

1995-11

A mathematical model for intra-cellular effects of toxins on DNA adduction and repair

Jacobs, Patricia A.

Monterey, California. Naval Postgraduate School

<http://hdl.handle.net/10945/24469>




Calhoun is a project of the Dudley Knox Library at NPS, furthering the precepts and goals of open government and government transparency. All information contained herein has been approved for release by the NPS Public Affairs Officer.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

NAVAL POSTGRADUATE SCHOOL

Monterey, California



**A MATHEMATICAL MODEL FOR
INTRA-CELLULAR EFFECTS OF TOXINS ON
DNA ADDUCTION AND REPAIR**

D. P. Gaver
P. A. Jacobs
R. L. Carpenter
J. G. Burkhart
November 1995

19951208 077

Approved for public release; distribution is unlimited.

Prepared for:

NPS Direct Funded Research Program

Naval Medical Research Institute/Toxicology Detachment
Wright-Patterson Air Force Base, Ohio

U.S. Army Biomedical Research & Development Laboratory
Ft. Detrick, MD 21702-5010

DTIC QUALITY INSPECTED 1

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943-5000

Rear Admiral M. J. Evans
Superintendent


Richard Elster
Provost

This report was prepared in conjunction with research funded by the Naval Medical Research Institute/Toxicology Detachment, Wright-Patterson Air Force Base, Ohio; the U.S. Army Biomedical Research & Development Laboratory, Ft. Detrick, MD 21702-5010; and the Naval Postgraduate School Direct Funded Research Program.

Reproduction of all or part of this report is authorized.

This report was prepared by:



DONALD P. GAVER, JR.
Professor of Operations Research


PATRICIA A. JACOBS
Professor of Operations Research



ROBERT L. CARPENTER
NMRI, Wright Patterson AFB, OH *OC*


JAMES G. BURKHART
NIEHS, Research Triangle Park, NC *JK*

Reviewed by:


FRANK PETHO
Acting Chairman
Department of Operations Research

Released by:


PAUL J. MARTO
Dean of Research

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 1995		3. REPORT TYPE AND DATES COVERED Technical
4. TITLE AND SUBTITLE A Mathematical Model for Intra-Cellular Effects of Toxins on DNA Adduction and Repair			5. FUNDING NUMBERS	
6. AUTHOR(S) Donald P. Gaver, Patricia A. Jacobs, Robert L. Carpenter, and James G. Burkhart				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School Monterey, CA 93943			8. PERFORMING ORGANIZATION REPORT NUMBER NPS-OR-95-012	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) NPS Direct Funded Research Program Naval Medical Research Institute/Toxicology Detachment Wright-Patterson Air Force Base, OH 45433-6503 U.S. Army Biomedical Research & Development Laboratory Ft. Detrick, MD 21702-5010			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The process by which certain classes of toxic compounds or their metabolites may react with DNA to alter the genetic information contained in subsequent generations of cells or organisms is a major component of hazard associated with exposure to chemicals in the environment. This paper formulates an <i>intra-cellular</i> dynamic model for one aspect of the action of toxins that form DNA adducts, when there is a capacity for removal of those adducts by a repair enzyme combined with reaction of the toxin and/or the DNA adduct to inactivate the repair enzyme. This particular model illustrates the possible saturation of repair enzyme capacity by the toxin dosage, and shows that bistable behavior can occur, with the potential to induce abrupt shifts between steady-state equilibria. The model suggests that bistable behavior, dose, and variation between individuals or tissues may combine under certain conditions to amplify the biological effect of dose observed as DNA adduction and its consequences as mutation. Models recognizing stochastic phenomena also indicate that variation in within-cell toxin concentration may promote jumps across stable equilibria.				
14. SUBJECT TERMS DNA adducts, DNA repair, mathematic models			15. NUMBER OF PAGES 36	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

D.P. Gaver^{1*}, P.A. Jacobs¹, R.L. Carpenter², and J.G. Burkhart^{3**}

The process by which certain classes of toxic compounds or their metabolites may react with DNA to alter the genetic information contained in subsequent generations of cells or organisms is a major component of hazard associated with exposure to chemicals in the environment. Many classes of chemicals may form DNA adducts and there may or may not be a defined mechanism to remove a particular adduct from DNA independent of replication. Many compounds and metabolites that bind DNA also readily bind existing proteins; some classes of toxins and DNA adducts have the capacity to inactivate a repair enzyme and divert the repair process competitively. This paper formulates an *intra-cellular* dynamic model for one aspect of the action of toxins that form DNA adducts when there is a capacity for removal of those adducts by a repair enzyme combined with reaction of the toxin and/or the DNA adduct to inactivate the repair enzyme. This particular model illustrates the possible saturation of repair enzyme capacity by the toxin dosage, and shows that bistable behavior can occur, with the potential to induce abrupt shifts away from steady-state equilibria. The model suggests that bistable behavior, dose, and variation between individuals or tissues may combine under certain conditions to amplify the biological effect of dose observed as DNA adduction and its consequences as mutation. Models recognizing stochastic phenomena also indicate that variation in within-cell toxin concentration may promote jumps across stable equilibria.

1. Department of Operations Research, Naval Postgraduate School, Monterey, CA 93943
2. Naval Medical Research Institute, Wright Patterson AFB, OH 45433
3. Environmental Toxicology Program, NIEHS, Research Triangle Park, NC 27709

**** Correspondent for biological issues**

Running Title: A mathematical model for removal of DNA adducts

3
rk, NC 27709

☒
☐
☐

Codes

Dist	Avail and/or Special
A-1	

1. Introduction

This paper formulates a simple generic mathematical model for one aspect of the action of toxins that bind DNA, react to inactivate a repair enzyme, and are removed within cells. It is an *intra-cellular* dynamic model that describes a process of formation of DNA adducts when there is a capacity for removal of those adducts from the DNA by a repair enzyme. Many compounds and metabolites that bind DNA also readily bind existing proteins; some classes of toxins and DNA adducts may inactivate the repair enzyme and divert the repair process competitively (Chae et al., 1994; Lijinsky et al., 1994; Mineura et al., 1994). This particular model illustrates the possible saturation of repair enzyme capacity by the toxin dosage, and shows that bistable behavior can sometimes occur, inducing abrupt shifts between two possible steady-state equilibrium. The bistable behavior, dose, and variation between individuals or tissues may combine under certain conditions to amplify the biological effect of dose, observed as DNA adduction. Models recognizing stochastic phenomena are also discussed; stochastic variation in within-cell toxin concentration can promote jumps across stable equilibria, as shown later.

The correct replication of a cell and its DNA is essential to the differentiation and maintenance of populations of cells within tissues and ultimately within host individuals, e.g. those making up a human population. Individual diploid human cells contain approximately 6×10^9 DNA base pairs comprising about 10^6 genes Burkhart (1995). Risk assessment technology and practice benefit from insights provided by biologically-based mechanistic models. The adduct/removal model presented here focuses only on one part of a complex intracellular process. Ultimately, models are required that describe the dynamics and behavior of dosimetry, cell surface recognition/transport, intracellular

DNA/protein interaction, repair, detoxification and clearance. At each step there is potential for a cell to control, modify, or possibly succumb to exogenously originating threats from toxic chemicals or their metabolites. Modification may also arise from interaction within cell populations through the action of exogenous signaling compounds. These interactions are to be the subjects of subsequent investigation.

2. Intra-Cellular Model of Damage and Repair

Chemical or radiological environmental stress may induce DNA damage as a *lesion*, such as a single or double DNA strand break, or oxidative- and hydroxyl radical- induced changes in structure. Damage also takes the form of an *adduct*, where reactive chemical groups may chemically react with the purine/ pyrimidine structures of DNA to disrupt hydrogen bond pairing or the action of polymerases (B. Singer, 1985; for review see *Drinking Water and Health.*). The adducted DNA, if not repaired by removal of the alkyl or aryl group may result in an altered DNA sequence (mutation) that is passed on to subsequent daughter cells. Depending on the site (gene) that is mutated, many biological outcomes are conceivable. If the event contributes to abnormal cell cycle regulation, such that the cell becomes a candidate for entry into the carcinogenic process, the interaction is of particular interest.

The most frequently described adducts have been methylated or ethylated nucleotides generated by direct alkylating agents (Bronstein et al., 1992; Pegg et al., 1990) These provide examples of the importance of the process in terms of mutation and the potential for biological variation between tissues and individuals to have profound effects on the mutagenic/carcinogenic outcome (Fox and Margison, 1988; Bronstein et al., 1991). Many toxic compounds are also likely to bind intra-cellular proteins, such as those involved in the removal of an

adduct from the DNA. There also may be a "suicide" reaction wherein the repair enzyme becomes inactivated and is thus removed from the population of molecules available for subsequent use by the cell in DNA repair (Hora et al., 1983; Pegg and Byers, 1992).

DNA adduction is one case among several possibilities, but it serves to motivate formulation of the generic mathematical models to follow. These are simplified extensively, but do capture some of the essentials of an adduct formation-repair/removal process in the presence of a toxic chemical and repair enzyme. Examination of the results of the model then suggests questions concerning system dynamics that are relevant to other intra- and extra-cellular processes, and to risk assessment. These must ultimately be answered by appropriate experiments.

2.1 Deterministic Mathematical Model

Suppose the cell in question is viewed at time t , where t may be measured from cell birth. To be specific let it be a stem cell, possibly in the spleen, cf. Alberts *et al.* (1994), so during time t the cell may have produced a number of daughter cells, while itself remaining alive.

We initially write a deterministic kinetic model to represent the system, utilize the following notation: $A(t)$ is the (mean) number of adducts present on the DNA of the cell, $R(t)$ is the (mean) number or concentration of *repair enzyme*, e.g. alkyl DNA transferase, present in the cell interior, and $T(t)$ is the concentration of the toxic chemical, or metabolite thereof, simultaneously present in the cell, all at time t . These quantities are stipulated to satisfy the following differential equations:

$$\frac{dA(t)}{dt} = \underbrace{\lambda_0 + \lambda_1 T(t)}_{\text{Adduct formation}} - \underbrace{\mu_{RA} R(t) A(t)}_{\substack{\text{Adduct removal/repair} \\ \text{by enzyme} \\ \text{(DNA transferase)}}} - \underbrace{\delta_{AM} A(t)}_{\substack{\text{Adduct removal} \\ \text{by mutation (plus} \\ \text{spontaneous disappearance)}}} \quad (2.1)$$

$$\frac{dT(t)}{dt} = \underbrace{\tau_C(t)}_{\substack{\text{Toxin input} \\ \text{to cell interior}}} - \underbrace{\delta_T T(t)}_{\substack{\text{Toxin removal;} \\ \text{other agents}}} - \underbrace{\mu_{RT} R(t) T(t)}_{\substack{\text{Toxin removal;} \\ \text{binds with} \\ \text{"suicide repair enzyme"} \\ \text{(DNA transferase)}}} \quad (2.2)$$

$$\frac{dR(t)}{dt} = \underbrace{\beta(\bar{R} - R(t))}_{\text{Enzyme creation}} - \underbrace{\delta_R R(t)}_{\substack{\text{Enzyme removal;} \\ \text{life-in-cell effect}}} - \underbrace{\mu_{RA} R(t) A(t)}_{\substack{\text{Enzyme removal} \\ \text{by adduct repair}}} - \underbrace{\mu_{RT} R(t) T(t)}_{\substack{\text{Enzyme removal;} \\ \text{binds with toxin}}} \quad (2.3)$$

These equations exhibit the possible double amplification effect of a toxin on adduct formation: first, the toxin contributes to the rate of adduct formation in accordance with rate parameter λ_1 , (see (2.1)); secondly, it is in competition for the repair enzyme, thus depleting the enzyme's level, in accordance with the rate parameter μ_{RT} (see (2.2)). The expression $\delta_T T(t)$ in 2.2 includes loss of toxin as a result first, of its removal during adduct formation as in (2.1); second, because of normal cellular processes such as detoxification or binding to DNA and proteins. Both a cell DNA adduct-enzyme product and a toxin-enzyme product may be induced; one example among many is the binding of benzylguanine to the alkyl DNA transferase (Chae et al., 1994).

The reader needs no reminder that the above model makes many simplifying assumptions, all of which may be questioned and modified as empirical information is developed. For example, the self-limiting enzyme creation term $\beta(\bar{R} - R(t))$ may be wrong in detail; even if adequate in mathematical form the upper limit, \bar{R} , may depend upon individual variation, tissue, cell age or cumulative exposure to toxin, possibly history-dependent expressed as an integral functional of $T(t')$, $0 \leq t' \leq t$. Similar comments may be made concerning

all of the rate parameters, λ_0 through μ_{RT} . For the present all parameters will be treated as constants. One aim of our modeling is to expose unexpected effects and sensitivities, and some such are revealed even for the present simplified setup. All assumptions made are hypothetical artifacts of the model until verified empirically, but may serve to suggest particular experiments.

3. Steady-State Solution

Suppose all parameters and the toxin input, τ_C , are *temporarily* assumed to be constants. If there is a long-run steady-state solution for $(A(t), T(t), R(t))$ then it satisfies the non-linear algebraic equations obtained by putting the derivatives equal to zero. In the present case steady-state solutions are obtained by solving the derivative = zero (2.1) for A in terms of R , (2.2) for T in terms of R , and finally putting these into (2.3). The result is

$$\begin{aligned} \beta \bar{R} = & (\beta + \delta_R)R + \mu_{RA} \left(\frac{\lambda_0}{\delta_{AM} + \mu_{RA}R} + \frac{\lambda_1 \tau_C}{(\delta_T + \mu_{RT}R)(\delta_{AM} + \mu_{RA}R)} \right) R \\ & + \frac{\mu_{RT} \tau_C R}{\delta_T + \mu_{RT}R}; \end{aligned} \quad (3.1)$$

after multiplying out by the denominator there results a cubic equation for the fixed points which are the appropriate solutions of (3.1). Although parametric solutions can be found for (3.1) it will not be easy to extract general qualitative information from them; see Appendix. A straightforward alternative is to introduce hypothetical biologically plausible parameters and to solve the differential equations (2.1), (2.2), and (2.3) numerically; this has been done in some trial cases.

Alternatively, qualitative information can be obtained from the formula (3.1), written as follows:

$$\begin{aligned} \beta(\bar{R} - R) = & \delta_R R + \left(\frac{\lambda_0 \mu_{RA}}{\delta_{AM} + \mu_{RA} R} + \frac{\tau_C \mu_{RT}}{\delta_T + \mu_{RT} R} \right) R \\ & + \frac{\lambda_1 \tau_C \mu_{RA} R}{(\delta_T + \mu_{RT} R)(\delta_{AM} + \mu_{RA} R)} \end{aligned} \quad (3.1a)$$

or

$$\ell(R) = r(R) \equiv r_1(R) + r_2(R) + r_3(R).$$

Graph the left-hand side of the above as a function of R , written $\ell(R) = \beta(\bar{R} - R)$

vs. the right-hand side, $r(R) = r_1(R) + r_2(R) + r_3(R)$ where $r_1(R) = \delta_R R$,

$$r_2(R) = \left(\frac{\lambda_0 \mu_{RA}}{\delta_{AM} + \mu_{RA} R} + \frac{\tau_C \mu_{RT}}{\delta_T + \mu_{RT} R} \right) R, \quad r_3(R) = \left(\frac{\lambda_1 \tau_C \mu_{RA} R}{(\delta_T + \mu_{RT} R)(\delta_{AM} + \mu_{RA} R)} \right).$$

Points of crossing are candidate solutions. Note the following possible qualitative configuration of the above components:




Figure 1

Now if the r.h.s. components are summed the following qualitative possibilities may emerge:




Figure 2A




Figure 2B




Figure 2C

The curve crossings, at values R_U and R_L in Figure 2, actually represent local equilibrium points; cf. Beltrami (1987). It is conjectured (and has been verified numerically in trial cases by examination of eigenvalues for the linearized system, using MATLAB) that in case the oscillating behavior of $r(R)$ occurs as shown, the value R_U in Fig. 2A, R_U and R_L in Fig. 2B, and R_L in Fig. 2C are all local stable points near which $R(t)$ will tend to reside as t increases, provided the starting level is near one of these values. If toxin dosage to the cell, τ_C , is low this is consistent with a relatively high level of enzyme, see formulas $r_2(R)$ and $r_3(R)$ in (3.1a) and Fig. 2A. If τ_C is high this is consistent with relatively low ambient enzyme: Fig. 2C. The intermediate equilibrium point in Fig. 2B is presumably always locally unstable. Fig. 2B suggests the existence of *two* such stable points, R_U and R_L ; if the enzyme level starts near one of these it tends to reside nearby, but if a disturbance (the origin of which is not modeled here) occurs, the enzyme level may abruptly shift, e.g. from R_U to R_L , consistent with a high level of ambient toxin *and* a relatively high formation rate for adducts. Switches back and forth could even occur. Such behavior has been called *bistable*. The conditions described tend to be associated with a relatively high rate of fixation by mutation when $R(t)$ is near R_L , with subsequent damaging effects. However, other protective behavior, such as apoptosis (programmed cell death), may be stimulated to inhibit overall adduct fixation and tumor production. Apoptosis is not modeled here.

3.1 Stochastic Model (Simulation)

The above deterministic model can be "made stochastic" in several ways, but the simplest, if not most elegant and complete, is to *computer-simulate*: (a) discretize time in equal-sized steps, (b) allow a discrete-time version of (2.1) to define the mean of a Markov stochastic process with (c) Normal/Gaussian

increments whose mean(s) are defined as in (b), and with variance equal to (or proportional to) the above incremental means, as would be appropriate for a diffusion approximation of a simple birth-death process. Formalize as follows, making time steps unity on an appropriate scale.

Mean Sequence

$$\begin{aligned} A(s+1) &= A(s) + \lambda_0 + \lambda_1 T(s) - \mu_{RA} R(s) A(s) - \delta_{AM} A(s) \\ &\equiv A(s) + \Delta A(s) \end{aligned} \quad (3.2)$$

$$\begin{aligned} T(s+1) &= T(s) + \tau_C(s) - \delta_T T(s) - \mu_{RT} R(s) T(s) \\ &\equiv T(s) + \Delta T(s) \end{aligned} \quad (3.3)$$

$$\begin{aligned} R(s+1) &= R(s) + \beta(\bar{R} - R(s)) - \delta_R R(s) - \mu_{RA} R(s) A(s) - \mu_{RT} R(s) T(s) \\ &\equiv R(s) + \Delta R(s) \end{aligned} \quad (3.4)$$

Stochastic Increments

$$\begin{aligned} \Delta A(s) &= \lambda_0(s) + \sigma_0(s) \Delta W_0(s) + \lambda_1 T(s) + \sigma_1(s) \Delta W_1(s) - \mu_{RA} R(s) A(s) \\ &\quad - \sigma_{RA}(s) \Delta W_{RA}(s) - \delta_{AM} A(s) - \sigma_{AM}(s) \Delta W(s) \end{aligned} \quad (3.5)$$

where

$$\begin{aligned} \sigma_0^2(s) &= \xi_0 \lambda_0(s) \\ \sigma_1^2(s) &= \xi_1 \lambda_1 T(s) \\ \sigma_{RA}^2 &= \xi_{RA} \mu_{RA} R(s) A(s) \\ \sigma_{AM}^2(s) &= \xi_{AM} \delta_{AM} A(s). \end{aligned} \quad (3.6)$$

The terms ΔW_i are mutually independent and Normal/Gaussian with mean 0 and variance 1 (if a different time step, the variance = time step). The constants $\xi_0, \xi_1, \xi_{RA}, \xi_{AM}$, are introduced so as to allow variability adjustment; putting them all equal to unity simulates a diffusion approximation to a simple birth-death model.

Likewise,

$$\begin{aligned}\Delta T(s) = & \tau_C(s) + \sigma_C(s)\Delta W_C(s) - \delta_T T(s) - \sigma_T(s)\Delta W_T(s) \\ & - \mu_{RT}R(s)T(s) - \sigma_{RT}(s)\Delta W_{RT}(s)\end{aligned}\quad (3.7)$$

$$\begin{aligned}\sigma_C^2(s) &= \xi_C \tau_C(s) \\ \sigma_T^2(s) &= \xi_T \delta_T(s)T(s)\end{aligned}\quad (3.8)$$

$$\sigma_{RT}^2(s) = \xi_{RT} \mu_{RT} R(s)T(s);$$

finally

$$\begin{aligned}\Delta R(s) = & \beta(\bar{R} - R(s)) + \sigma_\beta(s)\Delta W_\beta(s) - \delta_R R(s) - \sigma_R(s)\Delta W_R(s) \\ & - \mu_{RA}R(s)A(s) - \sigma_{RA}(s)\Delta W_{RA}(s) \\ & - \mu_{RT}R(s)T(s) - \sigma_{RT}(s)\Delta W_{RT}(s).\end{aligned}\quad (3.9)$$

It should be clear how to write down formulas for σ_β^2 and so forth.

Form the simulation

Realization

by recurrence:

$$A(s+1) = A(s) + \Delta A(s) \quad (3.10)$$

$$T(s+1) = T(s) + \Delta T(s) \quad (3.11)$$

$$R(s+1) = R(s) + \Delta R(s) \quad (3.12)$$

Start from initial conditions. Slight modifications will be necessary near boundaries to retain $R(s)$ -values positive and never greater than \bar{R} ; also $T(s)$ and $A(s)$ positive for all s . Note that there is a Normal/Gaussian Wiener process increment with each incremental component; some are common to two state-variable increments, e.g. ΔW_{RT} is in common with $\Delta T(s)$ and $\Delta R(s)$. The increment components that are not common to others can be combined so as to make necessary generation of just one (independent) increment component.

Mathematical Theory

Features of the escape from the neighborhood of either of two possible local stability points can be treated mathematically; this topic is called the *exit problem*, and is related to the *theory of large deviations*. Prominent contributors are Schuss (1980), Varadhan (1984), Aldous (1989), Simonian (1995), Freidlin and Wentzell (1984). An accessible textbook is Bucklew (1990). Attempts to mathematically calculate features of the process of jumping between stable points are not made in this paper; the qualitative features of the process are well-illustrated by simulation results.

Examples

Figure 3 through Figure 8 illustrate hypothetical time developments of adduct populations and corresponding amounts of enzyme. The initial values of A , R , and T are set equal to the largest root, R_U , of the cubic equation equivalent to (3.1) and the corresponding values of A and T ; the $\xi_C = 16$ and the other $\xi_i = 1$. Two replications are displayed in Figures 3 – 5; both depend on the same parameters, but flare-ups of adducts appear at quite different times; these high-adduct periods correspond to jumps from R_U to R_L (and in the present case rather quickly back), which are triggered by high fluctuations in internal toxin $T(t)$ caused initially by random fluctuations in $\tau(t)$, the amount of toxic chemical entering the cell. Figure 6 displays results for one replication. Figures 7 – 9 display the time series of the amount of enzyme, the number of adducts, and the amount of toxin along with histograms of the values. The histogram of the amount of enzyme has the most pronounced bimodal appearance. The lesser apparent bimodality in the histograms of number of adducts and the amount of toxin may reflect the fact that the derivatives of these two quantities are functions of the amount of enzyme and so the number of adducts and the amount of toxin

tend to be more related to the area under the enzyme time series and are thus smoothed. In these figures, all based on hypothetical parametric values, bistability is plainly visible. A burst of adducts can be expected to precede and trigger off other events upon successive replication after fixation of single or multiple mutations. Another possibility, also not currently modeled, is the possibility of actual cell death through necrosis or apoptosis when the intracellular toxin is relatively high and adducts are present.

4. Discussion

DNA abnormalities, e.g. adducts related to chemical exposure, are linked to mutation and the process of carcinogenesis. Hence their existence and relative prevalence within a tissue or cell may be useful as a risk analysis tool. However, the number of DNA adducts within any given cell of an organism on a temporal scale after acute or chronic exposures may be highly variable as a function of many interrelated biological processes.

In the present non-linear dynamic model the possible effect of initial conditions is dramatically illustrated: even a sudden brief shift of intra-cellular toxic chemical concentration can initiate a qualitative shift in cell condition, both inducing adduct formation and hobbling the cell's repair mechanism by binding the repair enzyme. The biological significance of the model is that such shifts, occurring within a context of replication across a number of cells can lead to adduct fixation as mutation and preconditions for transformation depending on the site, penetrance and expressivity of the event. The effect may be selective for individuals in the sense that some may have more resistance to assault owing to greater repair capability: larger enzyme production attributable to relatively larger values of β and \bar{R} in (2.3), perhaps leading to a greater-than-average repair enzyme concentration. Alternatively, there could exist a greater selectivity

in binding of the repair enzyme to DNA adducts rather than toxin, i.e. $\mu_{RA} > \mu_{RT}$ in (2.2) and (2.3) in some individual or tissues, thus leading to greater resistance to toxin. On the other hand, if propensity to form adducts is characteristic of at least some cells, i.e. reflected in relatively large parameters λ_0 and λ_1 , and perhaps also in outsized fixation rate δ_{AM} , see (2.1), then there can be inter-individual variations that could quantitatively describe an inherited predisposition to cancer as discussed by Sommerfeld, Meeker, Posadas, and Coffey (1995). This characteristic would be independent of the many preexisting mutations in genes known to be involved in the appearance of cancer in certain individuals.

REFERENCES

- Alberts, B., Bray, D., Lewis, J., Ratt, M., Roberts, K., and Watson, J.D., 1994. *Molecular Biology of the Cell*. Garland Publishing, Inc.
- Aldous, D., 1989. *Probability Approximations via the Poisson Clumping Heuristic*. Springer-Verlag, New York.
- Beltrami, E., 1987. *Mathematics for Dynamic Modeling*. Academic Press.
- Board on Environmental Studies and Toxicology. Commission on Life Sciences. National Research Council, 1989. Biologic significance of DNA adducts and protein adducts. *Drinking Water and Health*, Vol. 9: Selected Issues in Risk Assessment. National Academy Press, Washington DC, 6-59.
- Bronstein, S.M., Cochrane, J.E., Craft, T.R., Swenberg, J.A., and Skopek T.R., 1991. Toxicity, mutagenicity, and mutational spectra of *N*-ethyl-*N*-nitrosourea in human cell lines with different DNA repair phenotypes. *Cancer Res.* 51, 5188-5197.
- Bronstein, S.M., Skopek, T.R., and Swenberg, J.A., 1992. Efficient repair of *O*⁶-ethylguanine, but not *O*⁴-ethylthymine or *O*²-ethylthymine, is dependent upon *O*⁶-alkylguanine-DNA alkyltransferase and nucleotide excision repair activities in human cells. *Cancer Res* 52, 2008-2011.
- Bucklew, J.A., 1990. *Large Deviation Techniques in Decision, Simulation, and Estimation*. John Wiley & Sons, New York.
- Burkhardt, J.G., 1995. Perspectives on molecular assays for measuring mutation in humans and rodents. *Environ. Mol. Mutagen.* 25, 88-101.
- Chae, M.Y., McDougall, M.G., Dolan, M.E., Swenn, K., Pegg, A.E., and Moschel, R.C., 1994. Substituted *O*⁶-benzylguanine derivatives and their inactivation of human *O*⁶-alkylguanine-DNA alkyltransferase. *J. Med. Chem.* 37(3) 342-347.
- Fox, M. and Margison, G.P., 1988. Expression of an *E. coli* *O*⁶-alkylguanine DNA alkyltransferase gene in Chinese hamster cells protects against *N*-methyl and *N*-ethylnitrosourea induced reverse mutation at the hypoxanthine phosphoribosyl transferase locus. *Mutagenesis* 3, 409-413.
- Freidlin, M.I. and Wentzell, A.D., 1984. *Random Perturbations of Dynamical Systems*. Springer-Verlag, New York.
- Hodgman, C.D. (ed.) 1963. *CRC Standard Mathematical Tables*, Twelfth Edition, Chemical Rubber Publishing Company, Cleveland, OH.

- Hora, J.F., Eastman, A., Bresnick, E., 1983. O⁶-Methyl guanine methyltransferase rat liver. *Biochemistry* 22, 3759-3763.
- Lijinsky, W., Pegg, A.E., Anver, M.R., and Moschel, R.C., 1994. Effects of inhibition of O⁶-alkylguanine-DNA alkyltransferase in rats on carcinogenesis by methylnitrosourea and ethylnitrosourea. *Japan J. Cancer Res.* 85(3) 226-230.
- Michaelson, J., 1993. Biology of disease, cellular selection in the genesis of multicellular organization. *Laboratory Investigation*, 69, No. 2; 136-152.
- Mineura, K., Izumi, I., Watanabe, K., Kowada, M., Kohda, K., Koyama, K., Terashima, I., and Ikenaga, M., 1994. Enhancing effect of O⁶-alkylguanine derivatives on chloroethylnitrosourea cytotoxicity toward tumor cells. *Int. J. Cancer* 58(5) 706-712.
- Pegg, A.E., 1990. Mammalian O⁶ alkylguanine-DNA alkyltransferase: regulation and importance in response to alkylating carcinogenic and therapeutic agents. *Cancer Res* 50: 6119-6129.
- Pegg, A.E. and Byers, T.L., 1992. Repair of DNA containing O⁶-alkylguanine. *FASEB J.* 6(6) 2302-2310.
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., and Flannery, B.P., 1992. *Numerical Recipes in Fortran*, Second Edition, Cambridge University Press, Cambridge, UK.
- Schuss, Z., 1980. *Theory and Applications of Stochastic Differential Equations*. John Wiley & Sons, New York.
- Simonian, A., 1995. Asymptotic distribution of exit time for small-noise diffusions. *SIAM J. Applied Math.*, 55, 809-826.
- Singer, B., 1985. In vivo formation and persistence of modified nucleosides resulting from alkylating agents. *Environ. Health Persp.* 62, 41-48.
- Sommerfeld, H.J., Meeker, A.K., Posadas, E.M., and Coffey, D.S., 1995. Frontiers in prostate cancer, *CANCER Supplement*, Vol. 75, No. 7, 2027-2035.
- Strogatz, S.H., 1994. *Nonlinear Dynamics and Chaos*, Addison-Wesley Publishing Co.
- Varadhan, S., 1984. *Large Deviations and Applications*. SIAM, Philadelphia, PA.

APPENDIX

The Cubic Equation for Enzyme Fixed Points

The cubic equation that describes the fixed points of (2.1) – (2.3), specifically the solutions of (3.1), is as follows.

$$\begin{aligned}
 &(\beta + \delta_R)\mu_{RA}\mu_{RT}R^3 \\
 &+ [-\beta\bar{R}\mu_{RA}\mu_{RT} + (\beta + \delta_R)(\mu_{RA}\delta_T + \mu_{RT}\delta_{AM}) + \lambda_0\mu_{RA}\mu_{RT} + \tau_C\mu_{RT}\mu_{RA}]R^2 \\
 &+ [-\beta\bar{R}(\mu_{RA}\delta_T + \mu_{RT}\delta_{AM}) + (\beta + \delta_R)\delta_{AM}\delta_T + \lambda_0\mu_{RA}\delta_T + \lambda_1\tau_C\mu_{RA} + \tau_C\mu_{RT}\delta_{AM}]R \\
 &- \beta\bar{R}\delta_{AM}\delta_T = 0;
 \end{aligned} \tag{A.1}$$

division by $(\beta + \delta_R)\mu_{RA}\mu_{RT}$ puts this into the form in C.R.C. Standard Mathematical Tables or Press *et al.* (1992).

$$x^3 + ax^2 + bx + c = 0 \tag{A.2}$$

where

$$\begin{aligned}
 a &= \frac{[-\beta\bar{R}\mu_{RA}\mu_{RT} + (\beta + \delta_R)(\mu_{RA}\delta_T + \mu_{RT}\delta_{AM}) + (\lambda_0 + \tau_C)\mu_{RA}\mu_{RT}]}{(\beta + \delta_R)\mu_{RA}\mu_{RT}} \\
 b &= \frac{[-\beta\bar{R}(\mu_{RA}\delta_T + \mu_{RT}\delta_{AM}) + (\beta + \delta_R)\delta_{AM}\delta_T + \lambda_0\mu_{RA}\delta_T + \lambda_1\tau_C\mu_{RA} + \tau_C\mu_{RT}\delta_{AM}]}{(\beta + \delta_R)\mu_{RA}\mu_{RT}} \\
 c &= \frac{-\beta\bar{R}\delta_{AM}\delta_T}{(\beta + \delta_R)\mu_{RA}\mu_{RT}}.
 \end{aligned} \tag{A.3}$$

Next, put

$$Q = \frac{a^2 - 3b}{9}, \quad S = \frac{2a^3 - 9ab + 27c}{54}. \tag{A.4}$$

If $S^2 < Q^3$ there are three real roots; if $S^2 > Q^3$ there is one real root (and two imaginaries); if $S^2 = Q^3$ there will be equal roots.


No attempt has been made to simplify or interpret the formidable algebraic expressions that are obtained from the above formal manipulations. Note, however, that if there is no toxin input, i.e. $\tau_C(t) \equiv 0$, then $T(t) = 0$ and the cubic A reduces to the quadratic

$$(\beta + \delta_R)\mu_{RA}R^2 + [-\beta\bar{R}\mu_{RA} + (\beta + \delta_R)\delta_{AM} + \lambda_0\mu_{RA}]R - \beta\bar{R}\delta_{AM} = 0 \quad (\text{A.5})$$

It is apparent from the explicit solution that there will always be one positive fixed point, which will be the largest solution of (A.5). This will be a stable point. This conclusion follows directly from a graphical argument like that used before.

TWO REPLICATIONS INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20
SIG2C=25xSIG2C0



LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20





Figure 3

TWO REPLICATIONS INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20
SIG2C=25xSIG2C0



LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20





Figure 4

TWO REPLICATIONS INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20
SIG2C=25xSIG2C0



LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIME STEP=.01 BETA=4 RBAR=20




Figure 5

INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELI=.1

MURT=.6 DELR=.1

TIME STEP=.01 BETA=4 RBAR=20

XI(C)=25 OTHER XI=1







Figure 6

REPLICATION INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIMESTEP=.01 BETA=4 RBAR=20
XI(C)=25 OTHER XI(I)=1



HISTOGRAM OF AMOUNT OF ENZYME





Figure 7

REPLICATION INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1
MURT=.6 DELR=.1
TIMESTEP=.01 BETA=4 RBAR=20
XI(C)=25 OTHER XI(I)=1



HISTOGRAM OF NUMBER OF ADDUCTS




Figure 8


REPLICATION INITIAL ENZYME=17.14

LAM0=0 LAM1=10 MURA=.5 TAU=5 DELAM=.1 DELT=.1

MURT=.6 DELR=.1

TIMESTEP=.01 BETA=4 RBAR=20

XI(C)=25 OTHER XI(I)=1



HISTOGRAM OF AMOUNT OF TOXIN




Figure 9

DISTRIBUTION LIST

1.	Research Office (Code 08)	1
	Naval Postgraduate School Monterey, CA 93943-5000	
2.	Dudley Knox Library (Code 52)	2
	Naval Postgraduate School Monterey, CA 93943-5002	
3.	Defense Technical Information Center	2
	8725 John J. Kingman Rd., STE 0944 Ft. Belvoir, VA 22060-6218	
4.	Department of Operations Research	1
	Editorial Assistant (Code OR/Bi) Naval Postgraduate School Monterey, CA 93943-5000	
5.	Prof. Donald P. Gaver (Code OR/Gv)	5
	Naval Postgraduate School Monterey, CA 93943-5000	
6.	Prof. Patricia A. Jacobs (Code OR/Jc)	5
	Naval Postgraduate School Monterey, CA 93943-5000	
7.	Dr. J. Abrahams	1
	Code 111, Room 607 Mathematical Sciences Division, Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000	
8.	Dr. John Bailar	1
	468 N St. NW Washington, DC 20024	
9.	Prof. D. R. Barr	1
	Dept. of Systems Engineering U.S. Military Academy West Point, NY 10996	
10.	Dr. Frederic Bois	1
	IEP-MS90-3058 Lawrence Berkeley National Laboratory Berkeley, CA 94720	

11. Dr. David Brillinger.....1
 Statistics Dept.
 University of California
 Berkeley, CA 94720

12. Dr. James G. Burkhardt.....3
 MDC4-07
 Environmental Toxicology Program
 NIEHS
 Research Triangle Park, NC 27709

13. Prof. Brad Carlin.....1
 School of Public Health
 University of Minnesota
 Mayo Bldg. A460
 Minneapolis, MN 55455

14. Dr. Robert Carpenter.....10
 NAMRI/Navy Toxicology Detachment
 Bldg. 433, Area B
 Wright-Patterson AFB, OH 45433-6503

15. Center for Naval Analyses1
 4401 Ford Avenue
 Alexandria, VA 22302-0268

16. Prof. H. Chernoff1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

17. Mr. Harvey Clewell, III.....1
 ICF Kaiser Engineer
 ICF Kaiser International Inc.
 1201 Gaines Ave.
 Ruston, LA 71270-3107

18. Dr. Edward G. Coffman, Jr.1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974

19. Prof. John Copas1
 Dept. of Statistics
 University of Warwick
 Coventry CV4 7AL, ENGLAND

20. Prof. Sir David Cox1
 Nuffield College
 Oxford OX1 1NF, ENGLAND

21. Dr. Kenny S. Crump 1
Vice President, Environment Energy Group
ICF Kaiser Engineer
ICF Kaiser International Inc.
1201 Gaines Ave.
Ruston, LA 71270-3107

22. Prof. H. G. Daellenbach 1
Dept. of Operations Research
University of Canterbury
Christchurch, NEW ZEALAND

23. Dr. D. F. Daley 1
Statistics Dept. (I.A.S.)
Australian National University
Canberra, A.C.T 2606, AUSTRALIA

24. Dr. Naihua Duan 1
RAND Corporation
Santa Monica, CA 90406

25. Prof. Bradley Efron 1
Statistics Dept.
Sequoia Hall
Stanford University
Stanford, CA 94305

26. Dr. Guy Fayolle 1
I.N.R.I.A.
Dom de Voluceau-Rocquencourt
78150 Le Chesnay Cedex, FRANCE

27. Prof. George S. Fishman 1
Curr. in OR & Systems Analysis
University of North Carolina
Chapel Hill, NC 20742

28. Henry S. Gardner 1
U.S. Army Biological Research & Development Laboratory
Ft. Detrick, MD 21702-5010

29. Dr. Andrew Gelman 1
Statistics Dept.
University of California
Berkeley, CA 94720

30. Dr. Neil Gerr 1
Office of Naval Research
Arlington, VA 22217

31. Prof. Peter Glynn 1
Dept. of Operations Research
Stanford University
Stanford, CA 94305

32. Prof. Linda V. Green 1
Graduate School of Business
Columbia University
New York, NY 10027

33. Prof. J. Michael Harrison 1
Graduate School of Business
Stanford University
Stanford, CA 94305-5015

34. Dr. D. C. Hoaglin 1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138

35. Dr. David G. Hoel 1
Professor of Biometry and Epidemiology
Medical University of South Carolina
171 Ashley Ave.
Charleston, SC 29425-0002

36. Prof. D. L. Iglehart 1
Dept. of Operations Research
Stanford University
Stanford, CA 94305-5015

37. Institute for Defense Analysis 1
1800 North Beauregard
Alexandria, VA 22311

38. Dr. Robert C. Jackson 1
Vice President, Research and Development
Agouron Pharmaceuticals, Inc.
3565 General Atomics Court
San Diego, CA 92121-1121

39. Prof. J. B. Kadane 1
Dept. of Statistics
Carnegie-Mellon University
Pittsburgh, PA 15213

40. Dr. F. P. Kelly 1
Statistics Laboratory
16 Mill Lane
Cambridge, ENGLAND

41. Dr. Jon Kettenring 1
 Bellcore
 445 South Street
 Morris Township, NJ 07962-1910

42. Dr. Ralph Kodell 1
 Chief, Biometry Branch
 Biometry and Risk Assessment
 National Center for Toxicology Research
 3900 NCTR Drive
 Jefferson, AR 72079

43. Mr. Koh Peng Kong 1
 OA Branch, DSO
 Ministry of Defense
 Blk 29 Middlesex Road
 SINGAPORE 1024

44. Prof. Guy Latouche 1
 University Libre Bruxelles
 C.P. 212, Blvd. De Triomphe
 Bruxelles B-1050, BELGIUM

45. Dr. A. J. Lawrance 1
 Dept. of Mathematics
 University of Birmingham
 P.O. Box 363
 Birmingham B15 2TT, ENGLAND

46. Prof. M. Leadbetter 1
 Department of Statistics
 University of North Carolina
 Chapel Hill, NC 27514

47. Prof. J. Lehoczky 1
 Department of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

48. Dr. Georg Luebeck 1
 Fred Hutchinson Cancer Research Center
 1124 Columbia
 MP-665
 Seattle, WA 98014

49. Dr. Colin Mallows 1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974

50. Prof. R. Douglas Martin..... 1
Department of Statistics, GN-22
University of Washington
Seattle, WA 98195

51. Dr. Sati Mazumdar..... 1
Biostatistics Dept.
University of Pittsburgh
Graduate School of Public Health
Pittsburgh, PA 15261

52. Dr. James McKenna..... 1
Bell Communications Research
445 South Street
Morristown, NJ 07960-1910

53. Dr. Ramit Mehr-Grossman..... 1
Theoretical Biology and Biophysics
Theoretical Division
Mail Stop K710
Los Alamos National Laboratory
Los Alamos, NM 87545

54. Prof. Carl N. Morris..... 1
Statistics Department
Harvard University
1 Oxford Street
Cambridge, MA 02138

55. Dr. John A. Morrison 1
AT&T Bell Telephone Laboratories
600 Mountain Avenue
Murray Hill, NJ 07974

56. Prof. F. W. Mosteller 1
Department of Statistics
Harvard University
1 Oxford Street
Cambridge, MA 02138

57. Dr. John Orav 1
Biostatistics Department
Harvard School of Public Health
677 Huntington Avenue
Boston, MA 02115

58. Dr. Alan Perelson 1
Theoretical Biology and Biophysics
Theoretical Division
Mail Stop K710
Los Alamos National Laboratory
Los Alamos, NM 87545

59. Dr. Jim Petty 1
National Biological Survey
4200 New Haven Road
Columbia, MO 65201
60. Dr. Lorenz Rhomberg 1
Harvard Center for Risk Analysis
Harvard University
Cambridge, MA 02138
61. Dr. Rhonda Righter 1
Dept. of Decision & Info. Sciences
Santa Clara University
Santa Clara, CA 95118
62. Dr. John E. Rolph 1
Information and Operations Management
Univ. of Southern California
School of Business Administration
Los Angeles, CA 90089-1421
63. Prof. M. Rosenblatt 1
Department of Mathematics
University of California, San Diego
La Jolla, CA 92093
64. Prof. Frank Samaniego 1
Statistics Department
University of California
Davis, CA 95616
65. Prof. G. A. F. Seber 1
Dept. of Statistics
Univ. of Auckland
Private Bag 92019
Auckland, NEW ZEALAND
66. Prof. G. Shantikumar 1
The Management Science Group
School of Business Administration
University of California
Berkeley, CA 94720
67. Prof. N. D. Singpurwalla 1
George Washington University
Washington, DC 20052
68. Prof. H. Solomon 1
Department of Statistics
Sequoia Hall
Stanford University
Stanford, CA 94305

69. Dr. Andrew Solow 1
Woods Hole Oceanographic Institute
Woods Hole, MA 02543
70. Prof. W. Stuetzle 1
Department of Statistics
University of Washington
Seattle, WA 98195
71. Prof. J. R. Thompson 1
Dept. of Mathematical Science
Rice University
Houston, TX 77001
72. Prof. Steven K. Thompson 1
Statistics Dept.
Pennsylvania State Univ.
326 Classroom Bldg.
University Park, PA 16802-2111
73. Prof. J. W. Tukey 1
Statistics Dept., Fine Hall
Princeton University
Princeton, NJ 08540
74. Dr. D. Vere-Jones 1
Dept. of Math
Victoria Univ. of Wellington
P.O. Box 196
Wellington, NEW ZEALAND
75. Prof. David L. Wallace 1
Statistics Dept.
University of Chicago
5734 S. University Ave.
Chicago, IL 60637
76. Dr. Ed Wegman 1
George Mason University
Fairfax, VA 22030
77. Dr. L. Wein 1
Operations Research Center, Rm E40-164
Massachusetts Institute of Technology
Cambridge, MA 02139
78. Dr. Alan Weiss 1
Rm 2C-118
AT&T Bell Laboratories
600 Mountain Avenue
Murray Hill, NJ 07974-2040

79. Prof. Roy Welsch1
Sloan School
M.I.T.
Cambridge, MA 02139
80. Dr. Raymond S.H. Yang1
Colorado State University
College of Veterinary Medicine and Biomedical Sciences
Dept. of Environmental Health
Fort Collins, CO 80523